

# agKnowledge Spotlight

Shedding light on the concerns of your fields.



## Identifying Stalk Rot and Managing Potential Lodging Issues

- Lodging complicates corn harvest and can lead to yield loss and storage issues.
- Factors contributing to ear drooping and stalk lodging in corn include high winds, stalk cannibalization, and stalk rots.
- Understanding some of the contributing factors and how to most effectively manage the harvest and storage of lodged corn can assist in this year's harvest and planning for next season.

### Drooping Ears

Corn ears normally remain erect until after physiological maturity (black layer) when the shanks eventually collapse and the ears droop down (Figure 1). Recently, ears have begun to droop in drought-stressed fields that have not reached physiological maturity. Drooping ears suggest a loss of turgidity in the shank due to stress combined with cannibalization.

### Cannibalization and Physiological Stalk Lodging

In response to stress, corn plants will mobilize sugars to fill the kernels thus resulting in reduced sugar content of stalks. This process is referred to as stalk cannibalization and causes disintegration of the pith cells. The weakened stalks are more susceptible to colonization by fungi and to physiological stalk lodging. While this is often called stalk rot, the fungi are primarily colonizing tissues that are predisposed due to any condition that reduces photosynthesis and the production of carbohydrates needed to fill grain and maintain stalk integrity. Physiological stalk lodging is favored by good growing conditions early in the season, followed by stress after pollination. Stresses can include a lack of moisture, nitrogen deficiency, foliar disease, hail damage, and prolonged cool, cloudy weather conditions. Extended periods of dry or wet weather prior to pollination, followed by abrupt changes for several weeks after silking, also can cause poor stalk integrity and occurrence of physiological stalk lodging.

### Stalk Rots

Corn stalk rots are caused by fungi that are capable of invading otherwise healthy tissues. Stalk rots tend to be a complex of several disease-causing fungi. It is common for multiple causal organisms to be isolated from a single disease sample. Fields where stalk rot is developing should be identified and targeted for early harvest to minimize grain losses. The following are common stalk rots found in corn.

#### Anthracnose Stalk Rot

Symptoms usually occur just before the plant matures. Late in the season a shiny black discoloration develops as blotches or streaks on the stalk surface, especially on lower internodes (Figure 2).



Figure 1. A healthy corn ear (left) and a drooping corn ear (right). The milk line of these kernels are at 25 to 30% indicating several more days are needed prior to black layer.



Figure 2. Stalk rot symptoms. **Anthracnose** (left) - note the shiny, dark black blotches commonly seen on stalks. **Diplodia** (right) - symptoms include dark gray discoloration on stalk and the presence of pycnidia as indicated by the yellow arrow. This stalk is also exhibiting symptoms of anthracnose as noted by the red arrow.

Internal stalk tissue may become dark and soft, extending several internodes. Sometimes a portion of the plant above the ear dies prematurely (top dieback). More often, the entire plant is killed and several internodes are rotted. Lodging typically occurs higher on the stalk than with other stalk rots. Excellent resistance to anthracnose stalk rot is available in specific corn products.

#### Diplodia Stalk Rot

When a corn plant is infected by this disease, lower internodes become straw-brown, spongy, and dry. The pith tissues disintegrate, leaving vascular strands intact. White fungal mycelium may appear on the stalk surface. Tiny, dark fungal structures called pycnidia form just under the stalk surface (Figure 2). The pycnidia are embedded in outer stalk tissue and cannot be easily scraped off.

# Identifying Stalk Rot and Managing Potential Lodging Issues

## Fusarium Stalk Rot

Infection commonly leads to rotting of roots, crown, and lower internodes. Stalks prematurely dry down and are susceptible to breakage. When stalks are split, a whitish-pink to salmon discoloration may be visible. Disintegration of internal stalk tissues begins at the nodes (Figure 3). Crowns are frequently rotted.



**Figure 3.** Stalk rot symptoms. **Fusarium** (left) - typical symptoms are disintegration of internal tissues beginning at the nodes and rotted crown. **Gibberella** (right) - disintegration of pith tissues and pink to red discoloration.

## Gibberella Stalk Rot

Affected plants may wilt with leaves turning a dull gray-green. The lower stalk softens and becomes straw colored as plants die (Figure 3). Pith tissue disintegrates, leaving only vascular strands intact. The inside of a rotted stalk has a pink to red discoloration. Small, dark fungal bodies called perithecia form on the surface of the lower stalk. These perithecia are superficial and can be easily scraped off.

## Scouting

Fields should be scouted after pollination for visual symptoms of stalk rot or stalk cannibalization such as discoloration on stalks or early drydown. Different corn products and fields with different management practices should be evaluated separately.

When scouting, walk a zigzag pattern through the field and test stalk firmness by squeezing or pinching each stalk a couple of nodes above ground level. Healthy stalks are firm and cannot be compressed. If a stalk feels soft, it may be cannibalized and/or rotted and likely prone to lodging. Check at least 100 plants per field. If more than 10 to 15% of the stalks in a field are rotted, significant lodging is possible.<sup>2</sup>

A second method for determining potential stalk lodging is to randomly select 10 plants in a row and push each stalk 45 degrees (or about 5 to 8 inches) from upright. Repeat at 10 different locations within a field. If more than 10 to 15% of the stalks lodge or feel spongy, then the field should be slated for early harvest.<sup>3,4</sup>

## Harvesting Tips

Fields with considerable lodging or elevated lodging potential should be harvested early to help minimize the risk of further lodging and ear rots. Although drying cost is a concern when harvesting wet grain, this expense will likely be a better option compared to potential yield loss due to increased lodging and damage from ear rots. The following are some harvesting tips to protect yield potential:

- Corn reels can improve harvest efficiency.
- Harvest against the angle of the lodged corn to help maximize lift into the header.

- Harvesting when dew is present can minimize fluff.
- Combine should be adjusted to help minimize broken kernels and excess fines as they can lead to spoilage.
- Avoid over-threshing.
- Follow the combine manufacturer's manual for cylinder adjustments, speed, and clearance settings. Always refer to the manufacturer's manual before performing any maintenance.

## Grain Storage Tips

Once the grain is harvested, there is still a risk for loss. The following are some tips to help minimize grain storage losses:

- Wet corn in wagons or trucks should not be stored longer than 6 hours.
- Dry wet grain or put it in a holding bin for drying using forced air to keep it cool.
- Storing wet grain without aeration for 1 to 2 days can decrease storage life by 2 to 3 months.
- Mold growth can begin within 24 hours and accelerates rapidly if high-moisture corn is left in a wet bin too long.
- Check moisture content of every load and reset dryer controls based on changing moisture levels.
- Dry corn to 16% within 24 hours and cool to the outside air temperature within 48 hours.<sup>4</sup>
- Aerate to help corn attain a uniform temperature and to avoid "hot spots".
- Use stir augers to maintain airflow.
- Plan to check stored corn frequently. Stored corn should be inspected every 1 to 2 weeks in the fall and spring and once every 2 to 4 weeks after conditions in the bin have stabilized during the winter months.<sup>4</sup>

## Summary

Take the time to scout fields, regardless of when they were planted. Identify which fields may develop lodging issues and target these for an early harvest to help prevent potential harvest losses. When selecting corn products for next season it is important to note that ratings for stalk strength, lodging, and different diseases vary by product and can be influenced by genetic background, fertility levels, and management practices.

### Sources:

- 1 Corn stalk rots. 1995. University of Illinois Extension. RPD No. 200. <http://ipm.illinois.edu/>.
- 2 Munkvold, G. 2002. Time to start scouting for corn stalk rot. Integrated Crop Management.IC-488(20). Iowa State University Extension. <http://www.ipm.iastate.edu/ipm/icm/2002/8-19-2002/stalkrot.html>.<sup>3</sup> Vincelli, P. and Hershman, D.E. 1985. Corn stalk rots. PPA-26. University of Kentucky. [http://uknowledge.uky.edu/cgi/viewcontent.cgi?article=1045&context=anr\\_reports](http://uknowledge.uky.edu/cgi/viewcontent.cgi?article=1045&context=anr_reports).<sup>4</sup> McNeill S. and Montross, M. Corn harvesting, handling, drying, and storage. <http://www2.ca.uky.edu/>.<sup>5</sup> White, D. (editor). 1999. Compendium of corn diseases, Third edition. APS Press. Web sources verified 09/08/15.

For additional agronomic information, please contact your local seed representative. Developed in partnership with Technology, Development, & Agronomy by Monsanto.

Individual results may vary, and performance may vary from location to location and from year to year. This result may not be an indicator of results you may obtain as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible.

**ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** All other trademarks are the property of their respective owners. ©2015 Monsanto Company.  
130911070137 091015DLB